

# UNCONSTRAINED MINIMIZATION IN 1D

David F. Gleich

January 23, 2025

The material here is from Chapter 2 in Nocedal and Wright.

Everything will be 1d or univariate until further notice. Let  $f(x) : \mathbb{R} \rightarrow \mathbb{R}$ , for instance.

## 1 A BASKET OF DEFINITIONS

DEFINITION 1 (global minimizer) *A point  $x^*$  is a global minimizer if  $f(x^*) \leq f(x)$  for all  $x$  in the domain of  $f$ .*

DEFINITION 2 (strict global minimizer) *A point  $x^*$  is a strict global minimizer if  $f(x^*) < f(x)$  for all  $x$  in the domain of  $f$  except for  $x^*$*

DEFINITION 3 (local minimizer) *A point  $x^*$  is a local minimizer if  $f(x^*) \leq f(x)$  for all  $x$  in an open neighborhood of  $x^*$ .*

DEFINITION 4 (strict local minimizer) *A point  $x^*$  is a strict local minimizer if  $f(x^*) < f(x)$  for all  $x$  in an open neighborhood of  $x^*$  except for  $x^*$ .*

DEFINITION 5 (isolated minimizer) *A point  $x^*$  is an isolated (local) minimum if its a local minimizer and there is a neighborhood of  $x^*$  where  $x$  is the only minimizer.*

### 1.1 EQUIVALENCE?

Let  $f(x)$  be continuously differentiable. Can a strict minimizer be non-isolated?

*Isolated implies Strict*

*Strict implies Isolated?*

## 2 RECOGNIZING MINIMIZERS

*The key idea* Taylor's theorem ...

## 2.1 TAYLOR'S THEOREM (UNIVARIATE)

Let  $f(x)$  be twice continuously differentiable.

$$f(x_0 + h) =$$

$$f(x_0 + h) =$$

where  $\alpha$

## 2.2 NECESSARY CONDITIONS FOR A LOCAL MINIMIZER

*First order*

*Second order*

**Conclusion** So any local minimizer has:

## 2.3 SUFFICIENT CONDITIONS FOR A LOCAL MINIMIZER

*Necessary vs. Sufficient*

*Necessary* can be used to

*Sufficient* can be used to

*First order*

**To think about ...** What do these conditions say about  $f(x) = x^4$ ?